arXiv:1204.2397v1 [hep-ph] 11 Apr 2012

Collisional energy loss of heavy quarks

Alex Meistrenko[a], André Peshier[b], Jan Uphoff[a], Carsten Greiner[a]

[a]*Institut für Theoretische Physik, Johann Wolfgang Goethe-Universität Frankfurt, Max-von-Laue-Str. 1, D-60438 Frankfurt am Main, Germany*
[b]*University of Cape Town, Physics Department, Rondebosch 7701, South Africa*

Abstract

An analytic leading log formula for the averaged elastic energy loss of a heavy quark is found in the case of a non-static thermalized quark gluon plasma. Beyond the mean energy loss the collisional quenching of heavy quarks is considered in an improved pQCD transition matrix approach taking into account quantum distribution functions of medium particles, a running coupling and an effective screening mass adjusted to HTL calculations. The results for the nuclear modification factor and elliptic flow for RHIC energies within this model indicate a large contribution of the collisional energy loss of heavy quarks to their total quenching. Thereby hints for a qualitative difference between R_{AA} and v_2 were found, such that a linear scaling of the binary cross section cannot explain both effects at the same time. Therefore radiative energy loss and more precise initial conditions for the medium have to be taken into consideration.

Keywords: elastic energy loss, mean energy loss, heavy quarks, jet quenching, nuclear modification factor, elliptic flow

1. Introduction

Jet quenching, as observed in heavy ion collisions at RHIC [1, 2, 3, 4], has become an important probe of a deconfined state of matter with quasi-free quarks and gluons, the so-called quark gluon plasma (QGP). Indeed, this observable allows inferences on the properties of the expanding medium. Especially jet quenching of heavy flavors charm (c) and bottom (b) could be an ideal probe of the QGP. Due to the large mass[1] of these quarks, $m \gg T$, they are produced in a very early stage of the collision and can later be considered as test particles in the (equilibrated) heat bath.

The quenching of light quarks is well understood as an effect of radiative energy loss when propagating through the medium [5, 6]. However, in the case of heavy quarks pure radiative energy loss seems not sufficient to explain experimental data (mainly from electron yields) on heavy flavor quenching [7, 8, 9, 10], suggesting surprisingly strong attenuation. There is an ongoing debate on the reasons of quenching and energy loss of heavy partons in the QGP, which could be possibly clarified by analyzing and simulating spectra from heavy flavor decays. An additional contribution of suppression, significant in the range of small (to medium) momenta, might be the collisional energy loss by elastic processes as discussed in [10, 11, 12] and with further modifications in [13, 14, 15, 16, 17]. It will turn out that in our approach a fairly large collisional energy loss is predicted, although some comparisons between collisional and radiative contributions [18, 19] indicate another picture where the collisional loss should be relatively small ($\sim$ 20%) compared to the radiative one. These considerations are based on a fixed coupling and use a simple approximation for the screening mass in the QGP. In the following work both physical quantities are approximated in a more realistic way. Furthermore, references [13, 18] point out that some observables might be sensitive to the inclusion of binary collisions while the averaged energy loss only slightly increases by taking the collisional energy loss of parent partons into account, due to different probability distributions of radiative and collisional processes.

2. Mean energy loss with QGP flow

We start with a short overview of the mean energy loss for heavy quarks by binary collisions. In a static and thermalized plasma case (QED and QCD) a detailed recalculation of the energy loss per unit length dE/dx was done in [20, 21] to revise previous calculations by Braaten and Thoma [22, 23]. The general expression for the mean energy loss of a heavy quark with velocity $v = p/E$ and a scattering partner $i \in \{u, d, s, g\}$ from the medium is given by

$$\frac{dE_i}{dx} = \frac{1}{v} \int d\omega \, \frac{\partial \Gamma_i}{\partial \omega} \omega \,, \tag{1}$$

as the first moment of the differential transition rate $\partial\Gamma_i(E)/\partial\omega$ with respect to an energy transfer $\omega = E - E'$. If we are interested in computing the constant next to the leading logs of the energy loss (see (4)), we have to decompose the phase space of transfered momenta to a soft and a hard part by introducing $|t^*|$ as the separation scale, which can be chosen arbitrarily in the range of $m_D^2 \ll |t^*| \ll T^2$, where m_D^2 denotes the Debye screening mass and T the temperature of the system. There are two equivalent ways of computing transition rates either using self-energies or matrix elements [22]. That can be seen from cutting the one-loop self-energy diagram through the fermion

---

Email address: `meistren@uni-frankfurt.de` (Alex Meistrenko)

[1] $m_c \sim 1.3\,GeV$, $m_b \sim 4.6\,GeV$

lines. The soft part can be calculated in the so-called hard thermal loop (HTL) approximation from the self-energies as it was done in [20, 21]. The calculation of the hard contribution allows the second approach of computing interaction rates with tree-level matrix elements and using a bare propagator instead of a dressed one. Here the t-channel long-range gauge interactions are cut off by imposing a minimum momentum transfer $|t^*|$ and the expression for the mean energy loss reads:

$$\begin{aligned}\frac{dE_i}{dx} =& \frac{1}{2Ev}\int_k \frac{n_i(k)}{2k}\int_{k'}\frac{\overline{n_i}\,(k')}{2k'}\int_{p'}\frac{1}{2E'}\times\\ &\times(2\pi)^4\,\delta^{(4)}\left(P+K-P'-K'\right)\frac{1}{d}\sum_{spin,color}|\mathcal{M}_i|^2\,\omega\,,\end{aligned}\tag{2}$$

where $n_i(k)$ and $\overline{n_i}(k')$ in the expressions of the form $\int_k := \int d^3k/(2\pi)^3$ denote the thermal distribution functions. P and K stand for the initial and P', K' for the final four-momenta of the collisional partners (P, P' for the heavy quark). The sum of the matrix elements $|\mathcal{M}_i|^2$ runs over all initial and final states and is divided by the degeneracy factor $d = 6$ of the incoming heavy quark. In the limit of $E \gg T$ the equation (2) can be simplified to [20, 21]

$$\frac{dE_i}{dx} = d_i\int_k \frac{n_i(k)}{2k}\int_{t_{min}}^{t^*} dt(-t)\frac{d\sigma_i}{dt}\,,\tag{3}$$

here $d\sigma_i/dt$ are the Born cross sections [24] for the interaction with medium (target) particles of flavor i and degeneracy factor d_i. A rather long calculation of soft and hard phase space contributions leads to the following result for the mean energy loss (NLL-formula) of a heavy quark with mass M, which is independent of the separation scale $|t^*|$ [20, 21]:

$$\frac{dE}{dx} = \frac{4\pi\alpha_s^2T^2}{3}\left[\left(1+\frac{n_f}{6}\right)\ln\frac{ET}{m_D^2}+\frac{2}{9}\ln\frac{ET}{M^2}+c\left(n_f\right)\right].\tag{4}$$

Here $c(n_f)$ is a constant beyond the leading logarithms, which depends on the number of involved flavors $n_f = 3$ (u, d, s).

$$\begin{aligned}c(n_f) &= a\cdot n_f + b \simeq 0.146\cdot n_f + 0.050\\ a &= \left(\frac{2}{3}\right)\ln(2)-\frac{1}{8}+\frac{1}{6}c\\ b &= \left(\frac{31}{9}\right)\ln(2)-\frac{101}{108}+\frac{11}{9}c\\ c &= \frac{\zeta'(2)}{\zeta(2)}-\gamma\simeq -1.147\\ \gamma &\simeq 0.577\,\text{(Euler's constant)}\end{aligned}\tag{5}$$

This expression of the energy loss is valid for a fixed coupling α_s. A generalization to the running coupling can be found in [21]. As a remark, the first leading log arises from the t-channel interaction and was first anticipated by Bjorken [25].

In a numerical simulation we use the transition rate (see also next chapter) to determine the scattering probability and screen the t-channel by a mass term for regulating the infrared divergence. This was often done by using the Debye screening mass, which is not justified as can be seen in fig. 1, where the numerical result of the mean energy loss does not converge against the analytic formula (4). In fact, one has to introduce an effective screening mass of the form



Figure 1: Mean energy loss of a heavy quark as a function of the ingoing momentum for two different values of κ in comparison to the analytic NLL-formula (4).

$$\mu^2(t) = \kappa\cdot 4\pi\left(1+\frac{n_f}{6}\right)\alpha\,(t)\,T^2\,,\tag{6}$$

which can contain a fixed or running coupling $\alpha(t)$ and a free parameter κ. In the case of $\kappa = 1$ and a fixed coupling $\mu^2(t)$ reduces to the well-known (squared) Debye screening mass m_D^2. The procedure of introducing an effective screening mass is legitimate for the leading-order results because of the previous calculations (resumming and renormalizing) on the loop corrections for gluon exchange processes [26]. Therefore the coefficient κ can be calculated by evaluating the interaction rate $\partial\Gamma_i/\partial\omega$ of equation (1) with the effective cut-off μ^2 in the t-channel and comparing to the result of the NLL-formula (4). That leads to the following analytic relation for a fixed coupling [13, 14]

$$\kappa = \frac{1}{2e}\simeq 0.2\,.\tag{7}$$

With this value of κ we receive a convergence of the numerical results (our simulation is described in the next chapter) against the analytic formula (4) as expected, see fig. 1.

We next derive a relation for the leading log of the mean energy loss in the presence of a constant collective QGP flow. An interesting case with regard to a heavy ion collision is that of a parallel flow in respect to the propagation direction of a heavy quark, which interacts with the expanding medium. Here we expect a lower energy loss as predicted by the equation (4), whereas in the antiparallel case the heavy quark should slow down faster. As discussed before, the leading logarithmic accuracy does not require an HTL calculation, which will only contribute to a constant next to the leading log. Therefore it is sufficient to insert the Debye screening mass squared for the upper bound t_{max}, instead of a decomposition of the phase space. The thermal distribution functions $n_i(k)$ in the expression of the mean energy loss (2) have to be replaced by the Jüttner functions for bosons and fermions, which depend on the velocity of the medium $\vec{\beta}$ as well as Lorentz-factor γ (Lorentz boost of the four momentum):

$$n_{BJ} = \frac{1}{e^{\gamma\left(E_k - \vec{\beta}\cdot\vec{k}\right)/T} - 1}\,, \qquad n_{FJ} = \frac{1}{e^{\gamma\left(E_k - \vec{\beta}\cdot\vec{k}\right)/T} + 1}\,. \tag{8}$$

The lower bound on the integration variable t is set by the kinematics and can be approximated for $E \gg M^2/T$ in the following way:

$$t_{min} = -\frac{\left(s - M^2\right)^2}{s} \simeq -s \simeq -2Ek\,(1-x) \sim -O\,(ET)\;, \tag{9}$$

with

$$s = M^2 + 2PK + K^2 \simeq 2Ek\left(1 - \underbrace{\cos\vartheta}_{x:=}\right) \sim O\,(ET) \gg M^2\,, \tag{10}$$

where ϑ denotes the angle between $\vec{p}$ and $\vec{k}$. In the same high energy limit and $|t| \ll s$ for the leading log region of the t-channel, which is dominated by the infrared divergence $t \to 0$, we get

$$\begin{aligned} 2M^2 &= s + t + u \simeq s + u\,, \\ \Rightarrow \tilde{s} :&= \left(s - M^2\right) \approx \left(M^2 - u\right) =: -\tilde{u}\,. \end{aligned} \tag{11}$$

The evaluation of the t-channel integral with Born cross sections [24] and relations (9), (11) leads for quarks and gluons at leading log accuracy to

$$\begin{aligned} \sum_q \left.\frac{dE_q}{dx}\right|_t \simeq& \sum_q d_q \int_k \frac{n_{FJ}(k)}{2k} \int_{-2Ek(1-x)}^{-m_D^2} dt\,(-t)\,\times \\ &\times \frac{1}{16\pi\tilde{s}^2}\,\frac{64\pi^2\alpha_s^2}{9}\,\frac{\tilde{u}^2 + \tilde{s}^2}{t^2} \\ =& \frac{4\pi\alpha_s^2 T^2}{3}\,\frac{n_f}{6}\left[\ln\left(\frac{E\gamma\,(1-\beta)\,T}{m_D^2}\right) + const\right]\,, \end{aligned} \tag{12}$$

$$\begin{aligned} \left.\frac{dE_g}{dx}\right|_t \simeq& d_g \int_k \frac{n_{BJ}(k)}{2k} \int_{-2Ek(1-x)}^{-m_D^2} dt\,(-t)\,\times \\ &\times \frac{1}{16\pi\tilde{s}^2}\,32\pi^2\alpha_s^2\,\frac{-\tilde{s}\cdot\tilde{u}}{t^2} \\ =& \frac{4\pi\alpha_s^2 T^2}{3}\left[\ln\left(\frac{E\gamma\,(1-\beta)\,T}{m_D^2}\right) + const\right]\,. \end{aligned} \tag{13}$$

The u-channel calculation looks similar and requires only a change of variables from t to $\tilde{u}$. Here the dominant contribution to the cross section of the Compton scattering is proportional to $1/\tilde{u}^2$ and the relevant kinematic region is given by $-\tilde{s} = \tilde{u}_{min} \ll \tilde{u} \ll \tilde{u}_{max} = -M^2\tilde{s}/s$ as discussed in [20, 21]. The remaining matrix elements can be neglected, since they contribute only to the constant next to the leading log, thus

$$\begin{aligned} \left.\frac{dE_g}{dx}\right|_u \simeq& d_g \int_k \frac{n_{BJ}(k)}{2k} \int_{-2Ek(1-x)}^{-M^2} d\tilde{u}\,(\tilde{u} + \tilde{s})\,\times \\ &\times \frac{1}{16\pi\tilde{s}^2}\,\frac{64\pi^2\alpha_s^2}{9}\,\frac{-\tilde{s}\cdot\tilde{u}}{\tilde{u}^2} \\ =& \frac{4\pi\alpha_s^2 T^2}{3}\,\frac{2}{9}\left[\ln\left(\frac{E\,(\gamma\,(1-\beta))\,T}{M^2}\right) + const\right]\,. \end{aligned} \tag{14}$$



Figure 2: Mean energy loss for a jet of charm quarks $p_{jet} = 10\,GeV$ as a function of the medium velocity β according to the NLL formula (17) for different flow directions of the QGP with respect to the momentum vector of a charm quark.

So in the special case of a parallel flow ($\vec{\beta}\|\vec{p}$) we get

$$\begin{aligned} \left.\frac{dE}{dx}\right|_{\|}^{fix} = \frac{4\pi\alpha_s^2 T^2}{3}\Bigg[&\left(1 + \frac{n_f}{6}\right)\ln\frac{E\gamma\,(1-\beta)\,T}{m_D^2} \\ &+ \frac{2}{9}\ln\frac{E\gamma\,(1-\beta)\,T}{M^2} + c\left(n_f\right)\Bigg]\,, \end{aligned} \tag{15}$$

here the constant beyond logarithmic accuracy requires the tedious procedure of an HTL calculation but can be approximated by the constant $c(n_f)$ in the case of a static medium (5).

In the high energy limit $E \gg M$, which we use here, the term $E\gamma\,(1-\beta)$ can be replaced with the Lorentz invariant scalar product $p_\mu\beta^\mu$. That motivates a simple possibility of calculating the general form of dE/dx in the presence of an arbitrary but constant flow. Therefore we consider the rest frame (RF) of the QGP, where the heavy quark energy is determined by the general Lorentz boost of the four momentum vector to this frame. Using the equation (4), we get immediately the mean energy loss in the rest frame of the medium. Now we derive a relation for the differential energy loss between RF and a lab frame (LF), which is moving with a velocity $-\vec{\beta}$ relative to the rest frame. The only restriction in our argument is $p_\mu\beta^\mu \gg M$ at the lab frame. We start from the energy loss in RF (primed) and express all quantities by corresponding ones in LF (unprimed)

$$\begin{aligned} \frac{\Delta E'}{\Delta t'} = \frac{E_2' - E_1'}{t_2' - t_1'} &= \frac{\gamma\left(E_2 - \vec{\beta}\cdot\vec{p}_2\right) - \gamma\left(E_1 - \vec{\beta}\cdot\vec{p}_1\right)}{\gamma\left(t_2 - \vec{\beta}\cdot\vec{r}_2\right) - \gamma\left(t_1 - \vec{\beta}\cdot\vec{r}_1\right)} \\ &= \frac{\Delta E}{\Delta t}\,\frac{1 - \vec{\beta}\cdot\frac{\Delta\vec{p}}{\Delta E}}{1 - \vec{\beta}\cdot\frac{\Delta\vec{r}}{\Delta t}} \simeq \frac{\Delta E}{\Delta t}\,\frac{1 - \vec{\beta}\cdot\frac{\Delta\vec{p}}{\Delta p}}{1 - \vec{\beta}\cdot\frac{\Delta\vec{r}}{\Delta r}} = \frac{\Delta E}{\Delta t}\,\frac{1 - \vec{\beta}\cdot\frac{\Delta\vec{p}_\| + \Delta\vec{p}_\perp}{\Delta p}}{1 - \vec{\beta}\cdot\frac{\Delta\vec{r}_\| + \Delta\vec{r}_\perp}{\Delta r}} \\ &\simeq \frac{\Delta E}{\Delta t}\,\frac{1 - \vec{\beta}\cdot\frac{\Delta\vec{p}_\|}{\Delta p}}{1 - \vec{\beta}\cdot\frac{\Delta\vec{r}_\|}{\Delta r}} \simeq \frac{\Delta E}{\Delta t}\,\frac{1 - \beta\cos\sphericalangle\left(\vec{\beta}, \Delta\vec{p}_\|\right)}{1 - \beta\cos\sphericalangle\left(\vec{\beta}, \Delta\vec{r}_\|\right)} = \frac{\Delta E}{\Delta t}\,. \end{aligned} \tag{16}$$

Here, ⊥ and ∥ denote the transverse as well as parallel contributions relative to the initial momentum vector $\vec{p}_1$. The simple approximations used here are justified for the mean energy loss, since $\Delta\vec{p}/\Delta E \approx \Delta\vec{p}/\Delta p$ and $\Delta\vec{r}/\Delta t \approx \Delta\vec{r}/\Delta r$ are fulfilled for $p_\mu\beta^\mu \gg M$. Furthermore the scattering angle in the t- and u-channel for the same energy limit is negligibly small: $\Delta\vec{p}_\perp/\Delta p \approx \Delta\vec{r}_\perp/\Delta r \approx 0$. Thus the mean energy loss is approximately equal in both frames, that yields the leading logarithms in the lab frame without any complex calculations. We simply replace E of the equation (4) with the Lorentz invariant quantity $p_\mu\beta^\mu$ as long as the value of the product is much greater as the mass of the heavy quark. The constant next to the leading log would require a general evaluation of the integral (3) with Jüttner functions and an HTL calculation, which is not be investigated here. In general we obtain the following relation

$$\frac{dE}{dx} = \frac{4\pi\alpha_s^2T^2}{3}\left[\left(1+\frac{n_f}{6}\right)\ln\frac{p_\mu\beta^\mu T}{m_D^2} + \frac{2}{9}\ln\frac{p_\mu\beta^\mu T}{M^2} + c\left(n_f\right)\right]. \tag{17}$$

Fig. 2 illustrates the mean energy loss according to (17) for different flow directions ϑ and a fixed constant $c(n_f)$ to reproduce the prediction of the formula (4) at $\beta = 0$. We expect from the high energy behavior of the mean energy loss (consider (4), (16) and (17)) that the constant should be independent of the flow velocity $\vec{\beta}$.

3. MC-simulation with quantum transition rates

The simulation we use in the following is based on the double differential transition rate $\Gamma_2(p;p',\theta) := \partial^2\Gamma/\partial p'\partial\theta$ of binary collisions (Jacobian transformation of $\partial\Gamma_i/\partial\omega$ in (1)) in a thermalized medium of quarks and gluons. The distribution of Γ_2 is a function of the momenta p, p' and the scattering angle θ (see below). For our purpose Γ_2 is calculated by a numerical integration over finite size cells $\Delta p'\Delta\theta$. The divergence in the t-channel of the perturbative leading order (Born) cross sections is screened with an effective screening mass $\mu^2(t)$ as described before (6). For consistency and accurate predictions all α_s-dependencies in the screening mass and Born amplitudes are calculated with a running coupling as discussed in [13, 14]. This procedure generates a two dimensional matrix of transition rates. Multiplying the entries of the matrix with a small time step Δt leads to a probability matrix of a Markov chain [13]. The matrix contains all transition probabilities from an ingoing momentum p to another state with momentum p' and a scattering angle θ with respect to the direction of p. A set of such probability matrices for different ingoing momenta p form the basis of our MC-simulation for the collisional part of heavy quark suppression.

Fig. 3 shows a ratio of numerical to analytical results of the mean energy loss (17) with a fixed constant $c\left(n_f\right)$. This is actually a test of subleading terms, which were neglected in the calculation of (17). The figure underlines our prediction of the previous chapter that the mean energy loss is dominated by the leading log over a long range of β. Furthermore this comparison points out that the beyond leading log formula with flow produces reasonable results (deviation of $0-15\%$) even at rather small energies.



Figure 3: Comparison of numerical results for dE/dx to the analytical predictions for the NLL formula (17). The numerical calculations were done within the described MC-simulation for a jet of charm quarks with $p_{jet} = 10\,GeV$.

In our approach the heavy quarks are treated as test particles and do not affect the medium. This approximation is justified because of the small number of heavy flavor pairs, about 4 comparing to 800 gluons at mid-rapidity [27]. This allows us to use different backgrounds, such as hydro or BAMPS (Boltzmann Approach of Multi-Parton Scattering [28]), for the simulation of the local temperature in the medium, which determines the collisional transition rate of a thermalized plasma.

The spacial initialization of heavy quark dijet events is done according to the Glauber model [29, 30] and the initial momentum distributions are computed with the event generator *MC@NLO* for next-to-leading order calculations [31, 32]. Here the impact of shadowing and Cronin effect is neglected, since the modification of heavy quark distributions should be small for large momenta $p_T \gtrsim 2\,GeV$ [33]. An indirect comparison of initial momentum distributions to experimental data can be found in [17].

If at a later stage in our simulation the temperature of the medium drops below the critical temperature of $T_c \simeq 200\,GeV$, the heavy quarks pass a fragmentation process to heavy mesons. For this we use the Peterson fragmentation [34]

$$D_Q^M(z) = \frac{N}{z\left[1-\frac{1}{z}-\frac{\epsilon_Q}{1-z}\right]}, \tag{18}$$

where N is a normalization constant, $z = p_M/p_Q$ the momentum fraction (heavy meson to heavy quark) and $\epsilon_c = 0.05$, $\epsilon_b = 0.005$ typical squared mass ratios of involved quarks (light to heavy). The following stage of heavy meson decays to electrons is calculated with PYTHIA 8.1 [35, 36].

One important and free parameter in the described approach is the thermalization time τ, which has to be fixed by experimental data or theoretical models. It is not possible to define an exact value of the local temperature before equilibrium is reached and the calculation of scattering processes of heavy

	$T_0\,|\,T_c\,[GeV]$	$\tau_0\,|\,\tau_c\,[fm/c]$	$R\,[fm]$	dN_{ini}/dp_t^2
$set-1$	$0.42\,|\,0.2$	$0.6\,|\,5.6$	5.0	NLO
$set-2$	$0.30\,|\,0.2$	$1.0\,|\,3.4$	6.6	NLO

Table 1: Parameter sets for the Bjorken model, R denotes the effective radius of the nucleus.



Figure 4: Comparison of $c+b$, $D+B$ (Peterson fragmentation) and electron (decay from $D+B$ with PYTHIA) quenching at mid-rapidity in the Bjorken model to the observed R_{AA} from heavy flavor decays in the centrality class of 0% – 10% [7, 38].

quarks would not be very meaningful. This point will be discussed in the next section.

4. Heavy ion collisions

For transparency we first consider the particular steps of the quenching evolution (for quarks, mesons and electrons) in a simple model, namely within the Bjorken model, before we investigate more realistic scenarios. Furthermore we generalize the results of [13], because these were computed only for charm quarks in the transverse plane. Our observable is the nuclear modification factor, defined as a ratio of the final transverse momentum distribution to the distribution in a nucleon-nucleon collision, which is scaled with the number of binary collision $N_{b.coll.}$ according to the Glauber assumption

$$R_{AA} = \frac{d^2N_{AA}/dp_t dy}{N_{b.coll.} d^2N_{NN}/dp_t dy} \,. \tag{19}$$

We replace the power law momentum distributions of [13] by the NLO initial momentum distribution and consider the mid-rapidity for the same set of parameters (summarized in tab. 1), except the critical freeze-out temperature, which is set to $T_c = 200\,MeV$. That reduces the life time of the QGP phase $\tau_c = T_0^3\tau_0/T_c^3$ and ensures consistency to later results. Fig. 4 illustrates the combined modification factors of $c+b$, $D+B$ and electron yields for $set-1$ and $set-2$ (only e-data). From this plot it becomes apparent that the $c+b$ spectra are in good agreement with electron spectra from heavy flavor decays at large p_t-values ($p_t \geq 4\,GeV$), but slightly different at intermediate p_t and underestimate the depletion at smaller p_t. Our results for small momenta $p_t \lesssim 3\,GeV$ should be considered with some caution, as Peterson fragmentation is not suitable here and shadowing as well as Cronin effects were neglected. Even in such a simple framework we get hints for a significant contribution of the collisional energy loss with room for some radiative processes, hadronization and further effects. However the radial expansion will lead to a shorter life time of the plasma phase, therefore we will focus on more realistic dynamical models.

We can use available partonic or hydro models as input for the background medium by extracting the temperature and flow data of the discretized space for every time step. We focus here on the BAMPS medium, where the effective temperature $T_{cell}(\vec{x},t)$ and the flow velocity $\vec{v}_{cell}(\vec{x},t)$ are extracted locally. We then assume in each cell a complete thermodynamical equilibrium for the medium. In addition we also use a (3 + 1)-dimensional ideal hydrodynamic simulation (SHASTA [37]) as a control run to show that the observables are not really sensitive to differences in these models. Therefore we set the initial energy density of the ideal hydro to that of BAMPS by assuming a thermalization time of $\tau = 0.6 - 0.8\,fm/c$. These values of τ are in a good agreement with theoretical predictions and experimental data [28, 39] and are similar to the values of tab. 1. Fig. 5 confirms that the R_{AA} of charm quarks is not sensitive (neglect statistical fluctuations for $p_t \gtrsim 7\,GeV$) to a particular starting point of our simulation in the range of $t = 0.6 - 1.0\,fm/c$.

Figs. 6 and 7 present our results for the nuclear modification factor in a central and a peripheral collision with an impact parameter of $b = 8.2\,fm$, which corresponds to the centrality class of 20% – 40%. In both cases we get a quite good agreement with experimental data, if we keep in mind that the radiative energy loss is not taken into account. To mimic influence of radiative loss an artificial increase of the elastic cross section (transition rate) by a factor of $K = 3$ leads to an overestimation of the heavy quark quenching. In fact the experimental data is enveloped between $K = 1$ and $K = 3$ for high transverse momenta. Therefore we presume the total cross section of collisional and radiative processes to vary in this region. That means a smaller value of $K \approx 2$ in comparison to BAMPS results for R_{AA} and v_2 of heavy quarks, which need a $K \approx 4$ factor to be comparable with experimental data [17]. We emphasize that this observation can be explained from the difference between quantum versus Boltzmann distributed particles and from a difference of the number of active flavors (BAMPS does not consider light quarks in reference [17]). The formula of the mean energy loss (4) gets a prefactor of $\zeta(2)^{-1} = 6/\pi^2 \approx 0.61$ in the case of Boltzmann distributed medium particles, which gives a rough behavior for high jet energies.

After the nuclear modification factor R_{AA} we consider now the elliptic flow v_2, which is given by

$$v_2 = \left\langle \frac{p_x^2 - p_y^2}{p_t^2} \right\rangle . \tag{20}$$

The elliptic flow of heavy quarks forms at later stages of the collision by interactions with the medium. Consequently there is only a short and less important time period during the local



Figure 5: Sensitivity of R_{AA} in a peripheral nucleus-nucleus collision to different thermalization times $\tau = 0.6 - 1.0\,fm/c$, which are used as starting point of the simulation.

thermalization, which cannot be described correctly in our approach and is let out (no interactions for heavy quarks).

Fig. 8 shows v_2 in our model in comparison to the experimental data. Here we have to emphasize that an artificial increase of the cross section by a factor of $K = 3$ can only serve as an estimate of the experimental v_2 data. Comparing with the results of the full BAMPS cascade as published in [17], our factor for the K-value of 3 is on the same footing to the original prediction of BAMPS with $K = 4$.

5. Conclusion

We gave a brief overview of the derivation of the mean energy loss for a heavy quark in the QGP and generalized the formula to an arbitrary constant medium flow at leading log accuracy. These results were numerically tested using our transition matrix approach, which includes a running coupling as well as a modification of the Debye screening mass inferred by HTL considerations and is based on the transition rate for binary collisions in a thermalized medium of quarks and gluons.

In the second part of our work we reconsidered previous results on the elliptic flow and nuclear modification factor in the same framework of transition matrices. In the case of the nuclear modification factor an increase of the elastic cross section by a factor of $K < 3$ seems to reproduce experimental data of electrons from heavy flavor decays, whereas the elliptic flow requires $K \approx 3$. This discrepancy could be an effect of the radiative energy loss due to the non-isotropic gluon emission (see [40, 41]) and/or an impact of uncertain initial conditions, which could modify the transverse expansion of the medium.

As a final statement we emphasize that the elastic scattering alone cannot explain the sizeable nuclear modification factor and elliptic flow of heavy quarks but contributes significantly (20% – 50%) to their total quenching.



Figure 6: Nuclear modification factor R_{AA} at mid-rapidity $|y| < 0.35$ for a central $Au + Au$ collision with a center of mass energy of $\sqrt{s} = 200\,GeV$ (RHIC). The curves are calculated within our transition matrix approach for the NLO initial momentum distribution of heavy quarks. K denotes an artificial scaling factor of the elastic cross section. For comparison, experimental data on electrons from heavy flavor decays in the centrality class of 0% – 10% are shown [7, 38].



Figure 7: Same as fig. 6 but for a peripheral $Au + Au$ collision with an impact parameter of $b = 8.2\,fm$. For comparison, experimental data on electrons from heavy flavor decays in the centrality class of 20% – 40% are shown [38].



Figure 8: Same as fig. 7 but for elliptic flow.

Acknowledgments:

A. M. and J. U. acknowledge the kind hospitality at the University of Cape Town, where part of this work has been done. Part of the calculations was performed at the Center for Scientific Computing of the Goethe University Frankfurt. This work was supported by the Helmholtz International Center for FAIR within the framework of the LOEWE program launched by the State of Hesse.

References

[1] J. Adams *et al.* [STAR Collaboration], Nucl. Phys. A **757** (2005) 102 [arXiv:nucl-ex/0501009].
[2] K. Adcox *et al.* [PHENIX Collaboration], Nucl. Phys. A **757** (2005) 184 [arXiv:nucl-ex/0410003].
[3] I. Arsene *et al.* [BRAHMS Collaboration], Nucl. Phys. A **757** (2005) 1 [arXiv:nucl-ex/0410020].
[4] B. B. Back *et al.*, Nucl. Phys. A **757** (2005) 28 [arXiv:nucl-ex/0410022].
[5] R. Baier, Y. L. Dokshitzer, A. H. Mueller, S. Peigne and D. Schiff, Nucl. Phys. B **484** (1997) 265 [arXiv:hep-ph/9608322].
[6] M. Gyulassy, P. Levai and I. Vitev, Phys. Rev. Lett. **85** (2000) 5535 [arXiv:nucl-th/0005032].
[7] B. I. Abelev *et al.* [STAR Collaboration], Phys. Rev. Lett. **98** (2007) 192301 [Erratum-ibid. **106** (2011) 159902] [arXiv:nucl-ex/0607012].
[8] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **98** (2007) 172301 [arXiv:nucl-ex/0611018].
[9] J. Bielcik [STAR Collaboration], Nucl. Phys. A **774** (2006) 697 [arXiv:nucl-ex/0511005].
[10] S. Wicks, W. Horowitz, M. Djordjevic and M. Gyulassy, Nucl. Phys. A **784** (2007) 426 [arXiv:nucl-th/0512076].
[11] M. G. Mustafa and M. H. Thoma, Acta Phys. Hung. A **22** (2005) 93 [arXiv:hep-ph/0311168].
[12] G. D. Moore and D. Teaney, Phys. Rev. C **71** (2005) 064904 [arXiv:hep-ph/0412346].
[13] A. Peshier, arXiv:0801.0595 [hep-ph].
[14] P. B. Gossiaux and J. Aichelin, Phys. Rev. C **78** (2008) 014904 [arXiv:0802.2525 [hep-ph]].
[15] P. B. Gossiaux, R. Bierkandt and J. Aichelin, Phys. Rev. C **79** (2009) 044906 [arXiv:0901.0946 [hep-ph]].
[16] V. Greco, H. van Hees and R. Rapp, arXiv:1110.4138 [hep-ph].
[17] J. Uphoff, O. Fochler, Z. Xu and C. Greiner, Phys. Rev. C **84** (2011) 024908 [arXiv:1104.2295 [hep-ph]].
[18] G. -Y. Qin, J. Ruppert, C. Gale, S. Jeon, G. D. Moore and M. G. Mustafa, Phys. Rev. Lett. **100** (2008) 072301 [arXiv:0710.0605 [hep-ph]].
[19] B. G. Zakharov, JETP Lett. **86** (2007) 444 [arXiv:0708.0816 [hep-ph]].
[20] S. Peigne and A. Peshier, Phys. Rev. D **77** (2008) 014015 [arXiv:0710.1266 [hep-ph]].
[21] S. Peigne and A. Peshier, Phys. Rev. D **77** (2008) 114017 [arXiv:0802.4364 [hep-ph]].
[22] E. Braaten and M. H. Thoma, Phys. Rev. D **44** (1991) 1298.
[23] E. Braaten and M. H. Thoma, Phys. Rev. D **44** (1991) 2625.
[24] B. L. Combridge, Nucl. Phys. B **151** (1979) 429.
[25] J. D. Bjorken, FERMILAB-PUB-82-059-THY.
[26] A. Peshier, Phys. Rev. Lett. **97** (2006) 212301 [arXiv:hep-ph/0605294].
[27] J. Uphoff, O. Fochler, Z. Xu and C. Greiner, Phys. Rev. C **82** (2010) 044906 [arXiv:1003.4200 [hep-ph]].
[28] Z. Xu and C. Greiner, Phys. Rev. C **71** (2005) 064901 [hep-ph/0406278].
[29] R. J. Glauber, Nucl. Phys. A **774** (2006) 3 [nucl-th/0604021].
[30] M. L. Miller, K. Reygers, S. J. Sanders and P. Steinberg, Ann. Rev. Nucl. Part. Sci. **57** (2007) 205 [nucl-ex/0701025].
[31] S. Frixione and B. R. Webber, JHEP **0206** (2002) 029 [hep-ph/0204244].
[32] S. Frixione, P. Nason and B. R. Webber, JHEP **0308** (2003) 007 [hep-ph/0305252].
[33] N. Armesto, M. Cacciari, A. Dainese, C. A. Salgado and U. A. Wiedemann, Phys. Lett. B **637** (2006) 362 [hep-ph/0511257].
[34] C. Peterson, D. Schlatter, I. Schmitt and P. M. Zerwas, Phys. Rev. D **27** (1983) 105.
[35] T. Sjostrand, S. Mrenna and P. Z. Skands, JHEP **0605** (2006) 026 [arXiv:hep-ph/0603175].
[36] T. Sjostrand, S. Mrenna and P. Z. Skands, Comput. Phys. Commun. **178** (2008) 852 [arXiv:0710.3820 [hep-ph]].
[37] D. H. Rischke, S. Bernard and J. A. Maruhn, Nucl. Phys. A **595** (1995) 346 [nucl-th/9504018].
[38] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. C **84** (2011) 044905 [arXiv:1005.1627 [nucl-ex]].
[39] U. W. Heinz, AIP Conf. Proc. **739** (2005) 163 [nucl-th/0407067].
[40] R. Abir, C. Greiner, M. Martinez, M. G. Mustafa and J. Uphoff, arXiv:1109.5539 [hep-ph].
[41] R. Abir, U. Jamil, M. G. Mustafa and D. K. Srivastava, arXiv:1203.5221 [hep-ph].